

Recombinant Anti-CD98 Antibody (V3S-0522-YC5466)

Cat. No.: V3S-0522-YC5466

Summary

Description	This product is a monoclonal antibody derived from Human (<i>Homo sapiens</i>), which can specifically recognize Solute carrier family 3 member 2. The antibody is expressed with mammalian cell transient expression system, serum-free and purified by affinity chromatography. The purity and integrity are tested via SDS-PAGE and SEC-HPLC analysis. Given an antigen, additional QC measures are also desired such as affinity testing and binding validation. Specifically, the antibody is provided in multiple formats for in vivo and in vitro assays. The <i>In vivo</i> version features greater than 95% purity, ultra-low endotoxin levels (<1 EU/mg or 0.1 EU/mg), and is preservative, stabilizer, and carrier protein-free.
Clonality	Monoclonal
Host Species	Human
Target Species	Human
Immunogen	hCD98/hLAT1-E-expressing CT26 cell
Isotype	IgG, κ
Isotype Control	C34555
Secondary Antibody	C32400; C75370; C10513; C51635; C45530; C61612

Property

Expression Species	HEK293F or CHO
Conjugation	None
Purity	>95%, determined by SDS-PAGE and/or SEC-HPLC
Endotoxin	<1 EU/mg, determined by LAL method
Purification	Protein A affinity purified
Sterility	0.2 μ M filtered
Formulation	PBS, pH 7.4
Preservation	No preservatives
Stabilizer	No stabilizers
Storage	Store at 4°C within a week. For longer storage, aliquot and store at -20°C.

For lab research use only, not for diagnostic, therapeutic or any *in vivo* human use.

Applications

Application FC

Application Notes The antibody is recommended for detection of CD98 by FC assay.

Target

Target CD98

Alternative Name 4F2; CD98; MDU1; 4F2HC; 4T2HC; NACAE; CD98HC; SLC3A2

Gene ID [6520](#)

UniProt [P08195](#)

Introduction Solute carrier family 3 member 2 (SLC3A2, CD98 antigen) belongs to the solute carrier family and encodes a cell surface, transmembrane protein. The protein exists as the heavy chain of a heterodimer, covalently bound through di-sulfide bonds to one of several possible light chains. The encoded transporter plays a role in regulation of intracellular calcium levels and transports L-type amino acids.

Research Area Immunology; Signal Pathway

Related Disease Tumor

For lab research use only, not for diagnostic, therapeutic or any *in vivo* human use.